

38 - HIV Drug Resistance

Speaker: Michael Saag, MD

IDBR
INFECTIOUS DISEASE BOARD REVIEW
AUGUST 20-24
2022

HIV Drug Resistance

Michael S. Saag, MD
Professor of Medicine
University of Alabama at Birmingham




7/19/2022

IDBR
INFECTIOUS DISEASE BOARD REVIEW
AUGUST 20-24
2022

Disclosures of Financial Relationships with Relevant Commercial Interests





- None

How does resistance happen?




38 - HIV Drug Resistance

Speaker: Michael Saag, MD



Resistance Testing

- Genotypic resistance test
 - Perform test that gives mutations in viral genes
- Phenotypic resistance test
 - Perform test that describes growth of virus in the presence of anti-HIV drugs
- Limitations:
 - Cannot detect minority species (< 10% of viral population)



38 - HIV Drug Resistance

Speaker: Michael Saag, MD

Mutation Nomenclature

Codon (position)
PR = 1-99 amino acids
RT = 1-560 amino acids

M184V

Mutation Nomenclature

Codon (position)
PR = 1-99 amino acids
RT = 1-560 amino acids

M184V

Wild-type amino acid (consensus) → M
Mutant amino acid → V

Alanine	A
Cysteine	C
Aspartate	D
Glutamate	E
Phenylalanine	F
Glycine	G
Histidine	H
Isoleucine	I
Lysine	K
Leucine	L
Methionine	M
Asparagine	N
Proline	P
Glutamine	Q
Arginine	R
Serine	S
Threonine	T
Valine	V
Tryptophan	W
Tyrosine	Y

Mutations Selected by NNRTIs

Drug	Position	Wild-type	Mutations
Doravirine ¹²	106	A	V, G, P, F, M, L
	181	C, E, L	Y, G, P, F, M, L
Efavirenz	100, 101, 103, 106, 108	S, M, I	Y, G, P, F, M, L
	181	C, L, S	Y, G, P, F, M, L
Etravirine ¹³	50, 90, 100, 101, 106	S, M, I	Y, G, P, F, M, L
	181	C, L, S	Y, G, P, F, M, L
Nevirapine	100, 101, 103, 106, 108	S, M, I	Y, G, P, F, M, L
	181	C, L, S	Y, G, P, F, M, L
Raltegravir ¹⁴	100, 101	S, M	Y, G, P, F, M, L
	181	C, L, S	Y, G, P, F, M, L

Key Issues in HIV Resistance

- Easily Tested**
 - Specific Mutations
 - Cross – resistance
 - Prevalence of resistance at baseline
- Tough to Test**
 - Definition of Phenotypes
 - Complex resistance patterns
 - Genetic Barrier
 - Nuances of Resistance
 - Relationship between Pk and Pd

HIV Drug Resistance Testing

- Current guidelines recommend an **HIV genotype** as part of screening BEFORE ART is started.
- Following failure of 1st or 2nd regimens, **HIV genotype** is recommended to use with the history to choose the optimal next regimen.
- Following failure of 3rd and subsequent regimens, both **HIV genotype AND HIV phenotype** should be sent.
- If there is discordance between genotype and phenotype results, use the geno result (more sensitive).
- NOTE WELL:** Resistance mutations accrued from an earlier regimen **MAY NOT** be detected by tests obtained at the time of the current failing regimen

Everything You Need to Know About Nucleoside Analog Resistance in One Slide!

Mutation	Selected by	Effects on other NRTIs
184V	3TC, FTC	- Loss of susceptibility to 3TC, FTC - ↓ susceptibility to ABC, ddI (clinically insignificant) - Delayed TAMs and ↑ susceptibility to AZT, d4T, TDF
TAMs	AZT, d4T	- ↓ susceptibility to all NRTIs based on number of TAMs - More resistance with 41/210/215 than 67/70/219 pathway
151M, 69ins	AZT/ddI, ddI/d4T	- Resistance to all NRTIs - T69ins: TDF resistance
65R	TDF, ABC, ddI	- Variable ↓ susceptibility to TDF, ABC, ddI (and 3TC, FTC) - ↓ susceptibility to AZT
74V	ABC, ddI	- ↓ susceptibility to ABC, ddI - ↑ susceptibility to AZT, TDF
44D, 118I	AZT, d4T	- Increase NRTI resistance (with 41/210/215 pathway)

38 - HIV Drug Resistance

Speaker: Michael Saag, MD

CASE 1

- 25 year old man presents with newly diagnosed HIV
- Had an episode c/w acute seroconversion syndrome 4 months ago
- Initial HIV RNA 40,000; CD4 443 cells/ul
- He wants to start ARV therapy

Question #1

PREVIEW QUESTION

A baseline genotype is ordered that shows an M184V mutation. Which of the following drugs will have reduced susceptibility with this mutation?

- Efavirenz
- Zidovudine
- Tenofovir
- Etravirene
- Emtricitabine

DRUG	PHENOSENSE™ SUSCEPTIBILITY		Drug Sensitivity		NET ASSESSMENT
	Resistance	Drug Sensitivity	Drug	Pheno Score	
Abacavir	1.45	100%	ABC	Y	Sensitive
Didanosine	1.25	100%	ddl	Y	Sensitive
Emtricitabine	>MAX	100%	FTC	N	Reduced Susc.
Lamivudine	>MAX	100%	3TC	N	Reduced Susc.
Raltegravir	0.79	100%	ral	Y	Sensitive
Etravirene	0.27	100%	ZDV	Y	Sensitive
Zidovudine	0.45	100%	ZDV	Y	Sensitive
Tenofovir	0.45	100%	TDF	Y	Sensitive
NRTI Mutations: M184V					
Delamanvir	0.91	100%	DLV	Y	Sensitive
Etravirene	0.55	100%	EFV	Y	Sensitive
Nevirapine	0.53	100%	nef	Y	Sensitive
NRTI Mutations: none					

CASE 2

- 34 yo woman diagnosed with HIV 10 years ago
- Initially presented with PJP
- Initial Lab values
 - CD4 82 cells/uL
 - VL 106,000 c/mL
- Started on TDF / FTC / EFV (FDC)
- Did well for a while, then the regimen failed

Question #2

PREVIEW QUESTION

The genotype shows an M184V and K65R mutations. Which nRTI drugs would you include?

- ZDV
- TDF
- ddl
- ABC

Abacavir ¹⁴	E	L	V	Q
	65	74	115	104
	E	V	F	V
	N			
Emtricitabine	E			M
	65			104
	E			V
	N			I
Lamivudine	E			M
	65			104
	E			V
	N			I
Tenofovir ¹⁷	E	E		
	65	79		
	E			
	N			
Zidovudine ^{14,17}	M	D	E	L
	41	67	70	210,215,219
	L	N	R	W
				V
Didanosine ^{10,27}	E	L		
	65	74		
	E	V		
	N			

38 - HIV Drug Resistance

Speaker: Michael Saag, MD

Question #4

Which of the following results would indicate the highest likelihood of maraviroc activity?

- A. Pure R5 virus
- B. Pure X4 virus
- C. Mixture of R5 and X4 viruses
- D. Dual Tropic (R5/X4) virus

CASE 4


- 34 yo woman diagnosed with HIV 22 years ago
- Initially presented with PJP
- Initial Lab values
 - CD4 82 cells/uL
 - VL 106,000 c/mL
- Has been on multiple regimens over the years

Question #5


What is the likelihood she has high level resistance (< 2 active drugs available) ?

- A. < 1 %
- B. 1 - 5 %
- C. 5 -10%
- D. 10 - 20%
- E. > 20%

Prevalence of Patients with Limited Treatment Options



Virologic Success in Those with or without LTO




Common Mutations To Memorize

- M184V/I 3TC and FTC
- M41L, D67N, K70R, L210W, T215Y, K219Q "TAMS"
- 4 or more thymidine-analog mutations (TAMS) affect all approved nucleosides
- K65R tenofovir
- Q151M, 69SSS multi-NRTI
- K103N EFV (and NVP)
- retains susceptibility to etravirine
- Y181C NVP and other NNRTI
- E138K, K101E RPV and other NNRTI
- I50L ATV
- N155H, Q148H/R/K RAL and EVG
- Y143C RAL
- R263K DTG

38 - HIV Drug Resistance

Speaker: Michael Saag, MD



Summary

- High concern about resistance testing on Board Exams
- Difficult to create test questions that do not require complex interpretation, have a single best answer, or are not 'multiple true-false'
- Knowing common mutations and their role is a good way to prepare for the exam

37

