

09 - Penicillin Allergy

Speaker: Sandra Nelson, MD

2020

INFECTIOUS DISEASE BOARD REVIEW

Penicillin Allergy


Sandra B. Nelson, MD
Director, Musculoskeletal Infectious Diseases
Division of Infectious Diseases
Massachusetts General Hospital

Disclosures of Financial Relationships with Relevant Commercial Interests


None

Penicillin (PCN) Allergy: Premise

- 10% of the US population (32 million persons) have documentation of penicillin allergy
 - Rash (38%) > unknown (26%) > hives (18%) > angioedema (9%) > GI symptoms (6%) > itching (5%)
 - More common in older adults and hospitalized patients
- Vast majority of patients with penicillin allergy can be made to tolerate penicillin
 - Reactions are mild drug rashes that do not always recur
 - Reactions wane with time
 - Some reactions are not allergic




Penicillin Allergy



3 HARVARD

PCN Allergy: Consequences


- Alternative antimicrobial use
 - Less effective, more toxic, more broad spectrum
- Associated with:
 - increased risk of MRSA infections
 - increased risk of C difficile colitis
 - increased risk of surgical site infection
 - increased mortality
- An important target of stewardship efforts



4 HARVARD

Case #1

67 year old woman is hospitalized with nosocomial meningitis due to MSSA. She has a history of allergy to penicillin that is listed in the records as rash; the family recalls that she went to the ED when the rash occurred. She is not able to corroborate history. She has not received penicillin or cephalosporin antibiotics since the rash occurred a few years ago. Two of her daughters have allergies to penicillin.




5 HARVARD

Case #1: Vote

You are asked about optimal antibiotic treatment. What do you advise?

- A. Administer nafcillin without prior testing
- B. Administer nafcillin after test dose
- C. Skin test for penicillin reaction; if negative then administer nafcillin after test dose
- D. Administer vancomycin
- E. Desensitize to nafcillin




6 HARVARD

09 - Penicillin Allergy

Speaker: Sandra Nelson, MD

Options for Approaching PCN Allergy

- Monitored oral challenge
 - Useful for outpatients with low risk reactions (remote rash, pruritus) without imminent need of beta-lactam therapy
- Penicillin skin testing
 - Useful for inpatients and outpatients with a history of IgE mediated reaction, or sick patient with unknown reaction
 - Pre-Pen (benzylpenicilloyl polylysine) and penicillin G
 - Avoid in unstable patients



Shenoy JAMA 2019;321:188

Options for Approaching PCN Allergy


- Graded Challenge (1/10th test dose)
 - As a first step if suspicion for immediate reaction is low
 - After negative PCN skin testing when a related drug is desired (e.g. nafcillin)
- Desensitization
 - Positive skin test and/or confirmed immediate reaction, when a penicillin is the best therapy for an important infection
 - Desensitization wanes with missed doses (3 half-lives)
- Use of alternate therapy

Classification of Drug Allergy (Gell and Coombs)

Type	Immune mechanism	Clinical example	Management
I: Immediate (within one hour)	IgE-mediated	Anaphylaxis, Urticaria, Angioedema, Bronchospasm	Penicillin skin testing
II:	Antibody-dependent	Hemolytic Anemia, Thrombocytopenia, Neutropenia	No testing; generally avoid re-use
III	Immune Complex	Serum Sickness, Vasculitis	No testing; generally avoid re-use
IV delayed	Cell mediated	Cutaneous drug reactions, Interstitial nephritis, Hepatitis	Varies; for severe reactions and organ involvement avoid re-use

Deciphering Cutaneous Reactions

- IgE Mediated Reactions (hives)
 - Occur within minutes to hours
 - skin testing appropriate
 - if positive – desensitize or use alternate therapy
- Benign T-cell mediated
 - morbilliform or maculopapular
 - Usual onset days to weeks; persists >24 hours and resolves over days to weeks
 - test dose appropriate



Shenoy JAMA 2019;321:188

Deciphering Cutaneous Reactions

- Severe cutaneous reactions
 - DRESS and SJS/TEN
 - Usual onset days to weeks
 - Blistering, mucosal involvement, severe skin desquamation, organ involvement
 - avoid any beta-lactam
- Unknown Reaction
 - If hospitalized and/or critical illness:
 - Assume possibly IgE mediated
 - skin test then test dose



Stern NEJM 2012;366:2492

Shenoy JAMA 2019;321:188

Case #2

A 43 year old man with diabetes is hospitalized with a closed tibial fracture. Three years ago when he was being treated for a foot infection with piperacillin-tazobactam he developed a very itchy rash after several weeks of treatment. The anesthesiologist calls to ask advice about surgical antibiotic prophylaxis prior to operative fixation.

09 - Penicillin Allergy


Speaker: Sandra Nelson, MD

Case #2: Vote

- What do you do counsel?
- A. Administer clindamycin
 - B. Administer cefazolin
 - C. Administer cefazolin after intraoperative test dose
 - D. Administer ceftriaxone
 - E. Administer vancomycin


PCN Allergy and other beta-lactams

- Cephalosporins:
 - Significant cross reactivity 2%
 - Higher risk with earlier generation cephs
 - If suggestive type I PCN allergy:
 - use 3rd/4th gen (graded challenge preferred)
 - use 1st/2nd after PCN skin testing
 - If mild type IV reaction:
 - any cephalosporin OK
 - Avoid if severe reaction to PCN
- Carbapenems <1%
- Aztreonam: no cross reactivity



Cephalosporin Allergy

- Allergy often arises from side chains
 - More common than beta-lactam ring
- Probability of reaction higher when cephalosporins with similar side chains used ($R_1 > R_2$)
- Testable point:
 - Cefazolin has different side chains from all other cephalosporins



Thank you and good luck!



"The penicillin looks good."