

# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD



## Herpes Viruses: HSV and VZV in Immunocompetent and Immunosuppressed Patients

Richard J. Whitley, MD  
 Co-Director, Division of Pediatric Infectious Diseases  
 Loeb Eminent Scholar Chair in Pediatrics  
 Professor of Pediatrics, Microbiology, Medicine, and Neurosurgery  
 The University of Alabama at Birmingham

7/1/2024




### Disclosures of Financial Relationships with Relevant Commercial Interests

Steering Committee: NIAID COVID-19 Recover Study, NIAID Recover VITAL Study  
 Scientific Advisory Board: Treovir, LLC  
 Scientific Advisory Board: Altesa Biosciences  
 Member, Board of Directors at Evrys Bio  
 Member, Board of Directors at Virios Therapeutics  
 Past Chairperson: Merck Letemovir DMC, GSK IDMC for Zoster and NIAID COVID-19 Vaccine DSMB

## Herpes Viruses: The Family


- Herpes simplex virus, type 1 (HSV-1)
- Herpes simplex virus, type 2 (HSV-2)
- Varicella zoster virus (VZV)
- Cytomegalovirus (CMV)
- Epstein Barr virus (EBV)
- Human herpesvirus 6 (HHV 6 A and B)
- Human herpesvirus 7 (HHV 7)
- Human herpesvirus 8 (HHV 8)

## Viral Latency and Reactivation




Netter FH. ©2001 by Icon Learning Systems.

## Clinical Manifestations of Herpes Simplex Virus Infections



## Primary Herpes Simplex Virus Infection: Cutaneous Lesions



# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD

## Herpes Simplex Labialis




## Immunocompromised Host




## Most Widely Used Systemic Anti-HSV and VZV Drugs

- Acyclovir (ACV, Zovirax)
- Famciclovir (FCV, Famvir)
- Valacyclovir (VACV, Valtrex)
- Foscarnet (PFA, Foscavir)
- Ganciclovir (GCV, Cytovene)
- Val-Ganciclovir (Valcyte)
- Others:
  - Cidofovir



## Intravenous Acyclovir for Herpes Simplex Virus Infections in Immunocompromised Hosts

Time to cessation of viral shedding with acyclovir



## Acyclovir Prophylaxis for HSV Infection in BMT Patients

Acyclovir (250 mg iv/m2 /tid) or placebo for 18 days beginning 3 days before transplant

Group	Number of Patients	Number of HSV Infections	P
Acyclovir	10	0	~0.003
Placebo	10	7	

# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD




## Question #1



PREVIEW QUESTION

A 30 year old heart transplant has received acyclovir for the past 60 days for cutaneous HSV infection. The lesions are now progressive in spite of high-dose intravenous therapy. The most likely cause for disease progression is a deficiency or alteration of:

- A. Ribonucleotide reductase
- B. Reverse transcriptase
- C. Protease
- D. Thymidine kinase
- E. DNA polymerase



## Acyclovir Therapy of Genital Herpes

Summary of clinical benefit for treatment of:

- Primary
- Recurrent
- Suppressive


## Spectrum of HSV Clinical Presentation



First infection



Classical recurrence



Atypical recurrence

## Progression of Lesions



Early Redness/Swelling









Thin-Walled Fluid-Filled Vesicles and Pustules



Early Healing of Vesicles, Erosions, or Ulcers





# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD



# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD



- ### Genital Herpes: Viral Shedding
- Duration is longer in primary than in recurrent episodes
  - Higher rates in
    - People with frequent outbreaks
    - First year after acquisition
    - Primary: 12 days
    - Recurrent: 2-3 days
  - Oral antiviral suppressive therapy shortens the duration of, but does not eliminate, viral shedding
- Genital Herpes – A Clinician’s Guide to Diagnosis and Treatment. American Medical Association, 2001:1-20.  
Whitley RJ, et al. Clin Infect Dis. 1998;26:541-555.

### Herpes Presenting as Ulceration

- The patient had been to her doctor 3 times over the past 8 months with this pruritic and mildly painful rash on her right buttock. She had been told that it was an irritation from riding a bicycle.
- What is the key to the diagnosis?
  - A. the fact that lesions recurred
  - B. site of involvement is not unusual
  - C. trauma can induce reactivation

Photo courtesy of Jeffrey Gilbert, MD

# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD

## Question #2



PREVIEW QUESTION

An 18 year old man presents with a history of malaise, low-grade fevers, and new-onset painful genital lesions seen in the picture below. He had unprotected sexual intercourse with a female partner 2 weeks earlier. Neither he nor his partner has traveled outside the United States.



Which of the following diagnostic tests is most likely to yield the specific diagnosis?

- A. Serum RPR
- B. Serum FTA-Abs
- C. Darkfield microscopy
- D. Glycoprotein-G 1 serum antibodies
- E. PCR on lesion swab

## Oral Antiviral Therapies


- Famciclovir [Famvir®]
  - 500 mg
  - 250 mg
  - 125 mg
- Valaciclovir [Valtrex®]
  - 1 g
  - 500 mg
- Acyclovir [Zovirax®]
  - 800 mg
  - 600 mg
  - 200 mg

Valtrex® and Zovirax® are registered trademarks of GlaxoSmithKline.

## Impact of Acyclovir Therapy on Primary Genital HSV Infection

	Treatment Group (Days)		RR	P
	Acyclovir	Placebo		
Virus Shedding	2.8	16.8	6.82	0.0002
Pain	8.9	13.1	2.00	0.01
Scabbing	9.3	13.5	2.21	0.004
Healing	13.7	20.1	1.83	0.04

## Effect of Acyclovir Prophylaxis on Recurrent Genital Herpes



## Second Generation Anti-Herpetic Medications

- Valaciclovir (prodrug of acyclovir)
- Famciclovir (prodrug of penciclovir)



## Acyclovir/Valaciclovir Kinetics

DRUG	DOSE	PHARMACOKINETICS	
		C <sub>max</sub> (µg/mL)	Daily AUC (µg/mL·h)
VALTREX	1 g 3x/d	5.0	47
Oral ZOVIRAX	800 mg 5x/d	1.6	24
IV ZOVIRAX	5 mg/kg 3x/d	9.8	54
	10 mg/kg 3x/d	20.7	107


# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD

## Therapy of Recurrent Genital Herpes: Duration of Disease



## Famciclovir Therapy of Recurrent Genital Herpes



## Shorter and Shorter Therapy

- Genital Herpes
  - Valacyclovir: three days
  - Famciclovir: one day
- Labial Herpes
  - Valacyclovir: two days
  - Famciclovir: one day




## Prevention of Person-to-Person Transmission

## Valacyclovir Prevention of HSV Transmission to Susceptible Partners

Susceptible Partner	Val-ACV N = 743	Placebo N = 741	Total
No. acquired HSV-2	14	28	42
No. acquired HSV-1	0	4	4
No. developed clinical HSV-2	4	17	21

# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD



### HSE Morbidity


Percent Patients Patient Normal / Mild Impairment

Age	Glasgow Coma Scale	
	≤6	≥6
<30	0	60
>30	0	36

### Sensitivity and Specificity of PCR

	Biopsy Positive	Biopsy Negative
	PCR Positive	53
PCR Negative	1	44

Sensitivity 98%  
 Specificity 94%  
 Positive Predictive Value 95%  
 Negative Predictive Value 98%





# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts


Speaker: Richard Whitley, MD

## CHICKEN POX: Is Therapy of Value?

### Treatment of Chicken Pox: Adults (>18 Years) < 24 Hour Duration

	Acyclovir (n=38)	Placebo (n= 38)	P
Time to maximum number of skin lesions (days)	1.5	2.1	0.002
Days of new lesion information	2.7	3.3	0.03
Time to onset of cutaneous healing (days)	2.6	3.3	<0.001
Time to 100% crusting (days)	5.6	7.4	0.001
Maximum number of lesions	268	500	0.04

### Thoracic Herpes Zoster



### Questions

1. What is the most likely diagnosis?
2. How would you prove the etiology?



### Answer

- Clinically this is herpes zoster
- The lesion shown is Tzank prep positive on skin scraping. The sensitivity of this test is only ~60% and, therefore, is not recommended
- Immunofluorescence is positive for VZV, having a sensitivity of ~80%.
- Preferably, PCR can be performed even when lesions are scabbed and has the highest sensitivity.

### Question #3

What complication would you be most concerned about?

- A. Facial paralysis
- B. Keratitis
- C. Encephalitis
- D. Optic neuritis
- E. Oculomotor palsies



<http://www.ifnoroloji.org/kranialnoropatiler/Kranialnoropatiler.html>

## 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD

### Question #4 Stem

The patient has only the observed finding on his nose.

- What is your most likely diagnosis?
- What is the name of this sign?



www.medscape.com

### Question #4


What complication is it most likely to be associated with this illness?

- A. Deafness
- B. Vertigo
- C. Optic neuritis
- D. Keratitis
- E. Stroke

www.medscape.com

### Hutchinson's Sign

Zoster Involving nasociliary branch, Cranial Nerve VII which innervates the tip of the nose and the cornea



### Zoster Ophthalmicus



### NATURAL HISTORY OF ZOSTER IN THE NORMAL HOST

- Acute neuritis may precede rash by 48 - 72 hours
- Maculopapular eruption, followed by clusters of vesicles
- Unilateral dermatomal distribution

### NATURAL HISTORY OF ZOSTER IN THE NORMAL HOST

- Events of healing:
  - Cessation of new vesicle formation: 3 - 5 days
  - Total pustulation: 4 - 6 days
  - Total scabbing: 7 - 10 days
  - Complete healing: 2 - 4 weeks
- Cutaneous dissemination can occur  
dissemination is extremely rare
- Postherpetic neuralgia in 10% - 40% of cases

# 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD

## Complications of Zoster


### Common

- Postherpetic neuralgia
- Ocular complications
- Ophthalmic zoster
- (uveitis, keratitis, scleritis, optic neuritis)
- Pneumonitis
- Scarring
- Bacterial superinfection

### Uncommon

- Cutaneous dissemination
- Herpes gangrenosum
- Hepatitis
- Encephalitis
- Motor neuropathies
- Myelitis
- Hemiparesis (granulomatous CNS vasculitis)

## Prevalence and Duration of Pain




de Moragas et al. Arch Derm. 1957;75:193-196.


## Goals of Therapy

- Accelerate cutaneous healing
- Accelerate loss of pain acute / chronic
- Prevent complications

## Time to Cessation of Zoster-Associated Pain




## Time to Cessation of Zoster Associated Pain n = 1141



\* Beutner, et al. Acyclovir versus Valacyclovir in the treatment of herpes zoster in patients > 50 years old.

## Resolution of Pain in Herpes Zoster With Valacyclovir and Famciclovir



## 22 – HSV and VZV in Immuno-competent and Immunocompromised Hosts

Speaker: Richard Whitley, MD

### Summary of Efficacy of Concomitant Steroid Therapy with Acyclovir

- Accelerates resolution of acute neuritis
- Accelerates:
  - Return to usual activity P<0.001
  - Unaroused sleep P<0.0001
  - Cessation of analgesic use P<0.001
- Effect on chronic pain P=0.06

### Question #5

What is the most likely etiologic agent?



- A. HSV
- B. VZV
- C. CMV
- D. EBV
- E. HHV6

www.cdc.gov

### METHODS OF PREVENTING / MODIFYING VARICELLA

Pre-exposure: Oka varicella vaccine

Post-exposure: VZIG (now available in US)

Oka varicella vaccine  
(<3 days after exposure)  
Acyclovir  
(7-14 days after exposure)

### Shingles Prevention Trial: Zostavax

Attenuated, live virus (approved 2006)

- Efficacy but waning of immunity with time
  - Burden Of Illness 61.1% (51.1 – 69.1%)
  - Post-Herpetic Neuralgia 66.5% (47.5 – 79%)
  - Incidence of Herpes Zoster 51.3% (44.2 – 57.6%)

### Second Generation Vaccine: Shingrix

- Recombinant adjuvanted vaccine
  - Two shots
  - > 50 years of age
- Efficacy
  - Both PHN and incidence of shingles
  - >90% for >4 years
- Adverse events
  - Local reactogenicity: redness and pain ~ 50-70%
  - Systemic malaise/fever: ~30%

Thank You  
rwhitley@uab.edu