Speaker: John Bennett, MD



John E. Bennett, MD Bethesda, Maryland


7/1/2024



- Disclosures of Financial Relationships with Relevant Commercial Interests
  - None

## Mycology 101

- Yeasts reproduce by budding
  - All Candida have pseudohyphae in tissue except C.glabrata
  - Crypto has capsule, stains with mucicarmine
- Dimorphic fungi are round cells in tissue, hyphae in culture
  - Histoplasma, Coccidioides, Blastomyces, Sporothrix , Paracoccidioides
- Molds have hyphae in tissue and culture
  - Septate: Aspergillus, Fusarium, Scedosporium, others
  - Rare or no septae (Mucorales): Rhizopus, Mucor, Cunninghamella, others
  - Dark-walled fungi: many cause infection of skin, paranasal sinus, brain
     Phaeohyphomycosis




# Quick Quiz: What are these? Rhizopus calcofluor white Aspergillus wet mount Cladophialophora H&E

#### **ENDEMIC MYCOSES**

- Geographically restricted
- Dimorphic (yeast in tissue, hyphae in culture)
- •Infection by inhaling spores in nature
- •No person to person transmission
- Cluster of cases with fever, cough after soil exposure
  - No secondary cases
  - Desert dust=cocci. Rich earth, bat guano=histo
  - Streams, rivers=blasto

Speaker: John Bennett, MD




#### CASE 1

PREVIEW QUESTION

(COURTESY OF SHANAN IMMEL, MD)

- Formerly healthy 48M with 3 months of chronic fevers, cough, 25 Ib weight loss, night sweats, presented with acute worsening on dyspnea and was found to have a high fever and diffuse lung infiltrates bilaterally. Office worker in Md. No travel.Wife healthy.
- Vitals: 39.3C, HR 97, RR 29, BP 97/54, O2: 88% on room air
- · Crackle all over lung, spleen tip felt.
- WBC: 5,300, HgB 10.1 Plt 119,000, ALP 218, ALT 43, AST54, lactate 2.5, ferritin 2418, triglycerides 250. HIV neg.
- · Intubation, pressors, ceftriaxone, voriconazole





#### Recognizing mycoses on the board exam

#### Histoplasma capsulatum complex

- · Case clusters of acute pneumonia two weeks after soil exposure (rare: bat caves)
- Immunosuppressed patient with febrile disseminated disease


  - Cytopenias
    Miliary lung infiltrate can look like PIP, miliary TB
    Mucosal lesions resemble squamous carcinoma
    Adrenal insufficiency
    Can mimic HLH (hemophagocytic lymphohistiocytosis) or miliary TB
    HIV patients can have IRIS after starting ARV
- Urine or serum antigen good diagnostic test
- Biopsy: small budding yeast, mold on culture · Rx: ampho then itraconazole for disseminated
- Histoplasma duboisii (African histoplasmosis)






Speaker: John Bennett, MD














Speaker: John Bennett, MD





#### Case 3

The most likely fungus is which of the following:

- A. Scedosporium apiospermum (Pseudallescheria boydii)
- B. Lomentospora (Scedosporium) prolificans
- C. Apophysomyces elegans
- D. Fusarium multiforme
- E. Alternaria alternata

#### **Fusariosis**

Severely immunocompromised patients Mold, looks like Aspergillus in tissue Red, tender skin nodules

Routine blood culture grows mold in a third to half the patients

RX: response to amph and vori poor in severe neutropenia. Experimental: PMN transfusion?, fosmanogepix (investigational)??

Note: fungal meningitis from F. solani, Mexico, epidural anesthesia.

## Case 4

- 47 WM executive referred from Baltimore because of severe headaches, diplopia, high fever of 1 wk's duration
- 4 wks PTA: Maui resort one week
- 3 wks PTA: ranch outside Tucson, Arizona 1 wk
- 2 wks PTA: back at work in Baltimore
- 1 wk: PTA: Headache began
- Exam: Temp 38.5 C. Looks ill. Photophobia, nuchal rigidity, right CN6 palsy
- CBC, Routine blood chemistries normal. CSF: Glucose 55, Protein 58, WBC 330 (20% eos). Negative cryptococcal antigen on CSF, serum Lyme serology and serum RPR. MRI with contrast normal. Worsens during 2 wks of ceftriaxone. CSF cultures for bacteria, fungi, tbc neg to date.

#### CASE 4

The most helpful diagnostic test would be:

- A. CSF cytology
- B. Stool O&P
- C. Dietary history
- D. Fungal serology
- E. Leptospirosis serology

Speaker: John Bennett, MD

#### Coccidioidomycosis=Valley Fever

- Two species, one disease:
- C. immitis and C. posadasii. Both serious lab hazards Southwest USA. Washington state
- Acute pneumonia 2 wks after inhalation: arthralgias or erythema nodosum may accompany. Resolves.
- · Residual nodule or thin walled cavity may persist
- Dissemination: African americans, HIV, SOT, TNF inhibitors
- Bone, skin, chronic meningitis. Eosinophils
- Rx: fluconazole. Nonmeningeal: itraconazole

#### COCCIDIOIDOMYCOSIS DIAGNOSIS

#### SEROLOGY


CSF CF serology useful. Serum CF >16 suggests dissemination, falls with Rx Serum IgG by EIA converts to positive late, stays positive . Serum antigen in severe disease


#### CHITHE

Routine cultures negative, fungal cultures positive. Lab hazard

#### BIOPSY

Distinctive non-budding spherules





#### Blastomyces dermatitidis , B. gilchristii

CENTRAL USA AND CANADA, MOLD IN NATURE LARGE BROAD-BASED BUDDING IN TISSUE

MOIST EARTH NEAR RIVER, BEAVER DAMS.

NORMAL HOST


YEAST WITH BROAD BASED BUD, THICK WALL

ACUTE PNEUMONIA MAY SELF HEAL

INDOLENT, PROGRESSIVE PNEUMONIA DISSEMINATES TO SKIN, BONE, MALE GU TRACT

OFTEN PRESENTS AS SKIN LESIONS

RX: ITRACONAZOLE, AMPHO B





### CASE 6


Which is the most likely

- A. Babesia microti
- B. Candida tropicalis
- C. Fusarium oxysporum
- D. Aspergillus flavus
- E. Streptococcus anginosus

Speaker: John Bennett, MD

## Candidiasis makes the sick get sicker


- Fundoscopy for retinal lesions in candidemia patients.
   Intravitreal Rx may be needed
- · Remove intravenous catheter with candidemia
- Candida auris hospital outbreaks. Spreads on hands, surfaces
- Fluconazole resistance in C. auris, C. krusei, C. glabrata
- Fungitell (1-3) beta-D-glucan positive in serum



#### Case 7

32 yr old male with allogeneic hematopoietic stem cell transplant recipient for AML, developed graft versus host disease, given high dose prednisone, discharged and readmitted for fever not responding to antibacterial antibiotics. These two chest CT's, were taken at admission and a week later while he was responding to voriconazole. The most likely source of infection is:

- A. Dirt from his garden
- B. His oral floraC. Contaminated food
- D. Intravenous



### Aspergillus Pneumonia


Sudden onset of a <u>dense</u>, well circumscribed lesion in a neutropenic patient should suggest a mould pneumonia, most commonly aspergillosis, halo sign early, crescent sign later Septated hyphae invade blood vessels, infarct tissue.

Galactomannan useful in CSF, BAL, blood

False positives

False negatives with azole prophylaxis


Rx. voriconazole, isavuconazole, posaconazole, ampho B







Speaker: John Bennett, MD















Speaker: John Bennett, MD



#### MYCOSES WORTH MENTIONING

- SCEDOSPORIUM APIOSPERMUM: IMMUNOSUPPRESSED HOST CLINIALLY RESEMBLING ASPERGILLOSIS . BRAIN ABSCESS AFTER NEAR DROWNING IN POLLUTED WATER. AMPHOTERICIN B RESISTANT
- TRICHOSPORONOSIS: LIKE CANDIDIASIS BUT ECHINOCANDIN RESISTANT

